

(12) NACH DEM VERTRAG ÜBER DIE INTERNATIONALE ZUSAMMENARBEIT AUF DEM GEBIET DES
PATENTWESENS (PCT) VERÖFFENTLICHTE INTERNATIONALE ANMELDUNG

(19) Weltorganisation für geistiges Eigentum
Internationales Büro



(43) Internationales Veröffentlichungsdatum
3. Januar 2002 (03.01.2002)

PCT

(10) Internationale Veröffentlichungsnummer
WO 02/00928 A2

- (51) Internationale Patentklassifikation⁷: **C12Q 1/68** (81) Bestimmungsstaaten (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (21) Internationales Aktenzeichen: **PCT/EP01/07537**
- (22) Internationales Anmeldedatum:
2. Juli 2001 (02.07.2001)
- (25) Einreichungssprache: **Deutsch**
- (26) Veröffentlichungssprache: **Deutsch** (84) Bestimmungsstaaten (*regional*): ARIPO-Patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), eurasisches Patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), europäisches Patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI-Patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- (30) Angaben zur Priorität:
100 32 529.7 30. Juni 2000 (30.06.2000) DE
100 43 826.1 1. September 2000 (01.09.2000) DE
- (71) Anmelder (*für alle Bestimmungsstaaten mit Ausnahme von US*): **EPIGENOMICS AG [DE/DE]**; Kastanienallee 24, 10435 Berlin (DE).
- (72) Erfinder; und
- (75) Erfinder/Anmelder (*nur für US*): **OLEK, Alexander [DE/DE]**; Schröderstrasse 13/2, 10115 Berlin (DE). **PIEPENBROCK, Christian [DE/DE]**; Schwartzkopffstrasse 7b, 10115 Berlin (DE). **BERLIN, Kurt [DE/DE]**; Marienkäferweg 4, 14532 Stahnsdorf (DE).
- (74) Anwälte: **SCHOHE, Stefan** usw.; Boehmert & Boehmert, Franz-Joseph-Strasse 38, 80801 München (DE).

Veröffentlicht:

- ohne internationalen Recherchenbericht und erneut zu veröffentlichen nach Erhalt des Berichts
- mit dem Sequenzprotokollteil der Beschreibung in elektronischer Form getrennt veröffentlicht; auf Antrag vom Internationalen Büro erhältlich

Zur Erklärung der Zweibuchstaben-Codes und der anderen Abkürzungen wird auf die Erklärungen ("Guidance Notes on Codes and Abbreviations") am Anfang jeder regulären Ausgabe der PCT-Gazette verwiesen.

(54) Title: **DIAGNOSIS OF DISEASES ASSOCIATED WITH THE IMMUNE SYSTEM**

(54) Bezeichnung: **DIAGNOSE VON MIT DEM IMMUNSYSTEM ASSOZIIERTEN KRANKHEITEN**

(57) Abstract: The invention relates to chemically modified genomic sequences of genes associated with the immune system, an oligonucleotide directed against said sequence and/or PNA oligomers for the detection of the methylation state of cytosine of genes associated with the immune system. The invention also relates to a method for determining genetic and/or epigenetic parameters of genes associated with the immune system.

(57) Zusammenfassung: Die vorliegende Erfindung betrifft die chemisch modifizierte genomische Sequenzen von mit dem Immunsystem assoziierten Genen, gegen die Sequenz gerichtete Oligonukleotide und/oder PNA-Oligomere zur Detektion des Cytosin-Methylierungszustandes von mit dem Immunsystem assoziierten Genen sowie ein Verfahren zur Ermittlung von genetischen und/oder epigenetischen Parametern von mit dem Immunsystem assoziierten Genen.

WO 02/00928 A2

The remaining pages of this document are provided in electronic format on the attached compact disc.

Full paper copies will be provided upon request by the Examiner.

DIALOG(R)File 351:Derwent WPI
(c) 2002 Thomson Derwent. All rts. reserv.

014310206

WPI Acc No: 2002-130909/200217

Related WPI Acc No: 2001-596913; 2001-602751; 2001-602752; 2001-657177;
2002-010834; 2002-010922; 2002-010923; 2002-017444; 2002-017469;
2002-017470; 2002-017471; 2002-034446; 2002-090046; 2002-130908;
2002-139900; 2002-147896; 2002-154757; 2002-154758; 2002-154759;
2002-171649; 2002-351299; 2002-371829

XRAM Acc No: C02-040287

**Nucleic acid comprising fragment of chemically modified gene, useful for
diagnosis and treatment of diseases associated with abnormal cytosine
methylation**

Patent Assignee: EPIGENOMICS AG (EPIG-N)

Inventor: BERLIN K; OLEK A; PIEPENBROCK C

Number of Countries: 095 Number of Patents: 002

Patent Family:

Patent No	Kind	Date	Applicat No	Kind	Date	Week
WO 200200928	A2	20020103	WO 2001EP7537	A	20010702	200217 B
AU 200187575	A	20020108	AU 200187575	A	20010702	200235

Priority Applications (No Type Date): DE 1043826 A 20000901; DE 1032529 A
20000630

Patent Details:

Patent No	Kind	Lan	Pg	Main IPC	Filing Notes
-----------	------	-----	----	----------	--------------

WO 200200928	A2	G	32	C12Q-001/68	
--------------	----	---	----	-------------	--

Designated States (National): AE AG AL AM AT AU AZ BA BB BG BR BY BZ CA
CH CN CO CR CU CZ DE DK DM DZ EE ES FI GB GD GE GH GM HR HU ID IL IN IS
JP KE KG KP KR KZ LC LK LR LS LT LU LV MA MD MG MK MN MW MX MZ NO NZ PL
PT RO RU SD SE SG SI SK SL TJ TM TR TT TZ UA UG US UZ VN YU ZA ZW

Designated States (Regional): AT BE CH CY DE DK EA ES FI FR GB GH GM GR
IE IT KE LS LU MC MW MZ NL OA PT SD SE SL SZ TR TZ UG ZW

AU 200187575	A			C12Q-001/68	Based on patent WO 200200928
--------------	---	--	--	-------------	------------------------------

Abstract (Basic): WO 200200928 A2

NOVELTY - A nucleic acid (I) comprising a segment of at least 18
bases of a chemical pretreated DNA of a gene (A) associated with the
immune system, or its complement, where (A) is any one of 2420
sequences not given in the specification, is new.

DETAILED DESCRIPTION - INDEPENDENT CLAIMS are also included for the
following:

(1) oligomers (II) (oligonucleotides or peptide-nucleic acid (PNA)
oligomers) for detecting the status of cytosine methylation in
chemically treated DNA and comprising at least 9 nucleotides able to
hybridize to chemically pretreated (A) or their complements;

(2) a set of at least two (II);

(3) preparing an array of (II) on a solid phase;

(4) arrays produced by method (3);

(5) DNA and/or PNA arrays, for analysis of diseases related to
methylation status of genes, containing at least one (I);

(6) detecting genetic and/or epigenetic parameters for diagnosis
and/or treatment of diseases (or predisposition) by analysis of
cytosine methylation; and

(7) a kit containing bisulfite and (II).

ACTIVITY - Antiasthmatic; antiarteriosclerotic; antianemic;
cytostatic; nootropic; neuroprotective; anti-HIV; anticonvulsant;
ophthalmological; antirheumatic; antiarthritic; antidiabetic;
antipsoriatic; antiinflammatory. No details of tests for any of these
activities are given.

MECHANISM OF ACTION - None given in the source material.

USE - (I), and related oligomers (II) (or arrays of (II)), are
useful for diagnosis, prognosis and/or treatment of immune system

disorders, particularly where associated with aberrant cytosine methylation of the specified genes, e.g. eye diseases ((diabetic) retinopathy, neovascular glaucoma or macular degeneration); arteriosclerosis; anemia; (pancreatic) cancer; acute myeloid leukemia; Alzheimer's disease; AIDS; epilepsy; neurofibromatosis; rheumatoid arthritis; psoriasis; and inflammatory/ulcerative bowel diseases.

pp; 32 DwgNo 0/1

Technology Focus:

TECHNOLOGY FOCUS - BIOTECHNOLOGY - Preferred Gene: More than 300

(A) are tabulated, with their GenBank accession numbers.

Preferred Oligomer: These include at least one CpG dinucleotide, particularly in the middle third of the sequence. The set of at least 2 (II) particularly contains:

(i) (II) for determining the methylation status of all CpG dinucleotides from (A);

(ii) primer oligonucleotides for amplification of (A) or their complements (in this case the oligomers may be immobilized); or

(iii) at least 10 oligomers for detecting both cytosine methylation and/or single nucleotide polymorphisms in (A).

Preferred Array: (II) are arranged on a flat surface (of silicon, polystyrene, aluminum, steel, iron, copper, nickel, silver or gold) in a rectangular or hexagonal pattern.

Preferred Process: In method (6), a genomic DNA sample is treated, especially with bisulfite than alkali, to convert 5-unmethylated cytosines to uracil (or some other base with base-pairing properties different from those of cytosine) and fragments are amplified from the treated DNA using primer-(II). The amplicons, which include a detectable label, are then tested for hybridization to a set, or array, of probe-(II), and hybridized amplicons detected. At least 10 different fragments, of 100-2000 base pairs (bp), are amplified, in the same vessel, using a heat-stable polymerase and polymerase chain reaction for amplification. The amplicons are labeled with a fluorophore, radionuclide or a releasable molecular fragment of known mass, detectable by mass spectrometry (MS). Especially the amplicons, or their fragments (especially those with a single net charge for improved detection), are detected by MS, particularly using the matrix-assisted laser desorption/ionization or electrospray ionization techniques. The DNA being tested is isolated from e.g. cell lines, biopsies, blood, paraffin-embedded tissues etc.

